

Discussion 6A

The following notes are useful for this discussion: [Note 9](#), [Discussion 2A](#), [Homework 04](#)

1. Translating System of Differential Equations from Continuous Time to Discrete Time

Oftentimes, we wish to apply controls model on a computer. However, modeling a continuous time system on a computer is a nontrivial problem. Hence, we turn to discretizing our controls problem. That is, we define a discretized state $\vec{x}_d[i]$ and a discretized input $\vec{u}_d[i]$ that we “sample” every Δ seconds. The notion of discretization is very similar to the approach covered in [Discussion 2A](#).

(a) Consider the scalar system below:

$$\frac{dx(t)}{dt} = \lambda x(t) + bu(t). \quad (1)$$

where $x(t)$ is our state and $u(t)$ is our control input. Let $\lambda \neq 0$ be an arbitrary constant. Further suppose that our input $u(t)$ is piecewise constant, and that $x(t)$ is differentiable everywhere (and thus, continuous everywhere). That is, we define an interval $t \in [i\Delta, (i+1)\Delta)$ such that $u(t)$ is constant over this interval. Mathematically, we write this as

$$u(t) = u(i\Delta) = u_d[i] \text{ if } t \in [i\Delta, (i+1)\Delta). \quad (2)$$

The now-discretized input $u_d[i]$ is the same as the original input where we only “observe” a change in $u(t)$ every Δ seconds. Similarly, for $x(t)$,

$$x(t) = x(i\Delta) = x_d[i] \quad (3)$$

Let’s revisit the solution for eq. (1), when we’re given the initial conditions at t_0 , i.e we know the value of $x(t_0)$ and want to solve for $x(t)$ at any time $t \geq t_0$:

$$x(t) = e^{\lambda(t-t_0)}x(t_0) + b \int_{t_0}^t u(\theta)e^{\lambda(t-\theta)} d\theta \quad (4)$$

Given that we start at $t = i\Delta$, where $x(t) = x_d[i]$ is known, and satisfy eq. (1), where do we end up at $x_d[i+1]$? (HINT): Think about the initial condition here. Where does our solution “start”?

Solution: For $t \in [i\Delta, (i+1)\Delta)$, the differential equation takes the form

$$\frac{dx(t)}{dt} = \lambda x(t) + bu(t) = \lambda x(t) + bu_d[i] \quad (5)$$

where we choose our initial condition to be $x(i\Delta) = x_d[i]$, since this is a known quantity. We can solve this equation for $x(t)$ using the integral equation from eq. (4) and the fact that $u_d[i]$ is a constant value over this interval. In particular, we get the following form

$$x(t) = e^{\lambda(t-i\Delta)} \underbrace{x(i\Delta)}_{x_d[i]} + b \int_{i\Delta}^t \underbrace{u(i\Delta)}_{u_d[i]} e^{\lambda(t-\theta)} d\theta \quad (6)$$

$$= e^{\lambda(t-i\Delta)} x_d[i] + bu_d[i] \int_{i\Delta}^t e^{\lambda(t-\theta)} d\theta \quad (7)$$

Plugging in the timestep of interest, we set $t = (i+1)\Delta$, to evaluate $x_d[i+1]$ as

$$x_d[i+1] = x((i+1)\Delta) \quad (8)$$

$$= e^{\lambda\Delta} x_d[i] + bu_d[i] \int_{i\Delta}^{(i+1)\Delta} e^{\lambda((i+1)\Delta-\theta)} d\theta \quad (9)$$

$$= e^{\lambda\Delta} x_d[i] + bu_d[i] \frac{e^{\lambda\Delta} - e^0}{\lambda} \quad (10)$$

$$= e^{\lambda\Delta} x_d[i] + bu_d[i] \frac{e^{\lambda\Delta} - 1}{\lambda} \quad (11)$$

which gives us the solution for $x_d[i + 1]$.

- (b) Suppose we now have a continuous-time system of differential equations, that forms a vector differential equation. We express this with an input in vector form:

$$\frac{d\vec{x}(t)}{dt} = A\vec{x}(t) + \vec{b}u(t) \quad (12)$$

where $\vec{x}(t)$ is n -dimensional. Suppose further that the matrix A has distinct and non-zero eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_n$, with corresponding eigenvectors $\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n$. We collect the eigenvectors together and form the matrix $V = [\vec{v}_1, \vec{v}_2, \dots, \vec{v}_n]$.

We now wish to find a matrix A_d and a vector \vec{b}_d such that

$$\vec{x}_d[i + 1] = A_d \vec{x}_d[i] + \vec{b}_d u_d[i] \quad (13)$$

where $\vec{x}_d[i] = \vec{x}(i\Delta)$.

Firstly, define terms

$$e^{\Lambda\Delta} = \begin{bmatrix} e^{\lambda_1\Delta} & 0 & \dots & 0 \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & e^{\lambda_n\Delta} \end{bmatrix} \quad (14)$$

$$\Lambda^{-1} = \begin{bmatrix} \frac{1}{\lambda_1} & 0 & \dots & 0 \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & \frac{1}{\lambda_n} \end{bmatrix} \quad (15)$$

$$\vec{u}_d[i] = V^{-1} \vec{b} u_d[i] \quad (16)$$

Note that the term $e^{\Lambda\Delta}$ is just a label for our intents and purposes — this is not the same as applying e^x to every element in the matrix Λ .

Complete the following steps to derive a discretized system:

- i. **Diagonalize the continuous time system using a change of variables (change of basis) to achieve a new system for $\vec{y}(t)$.**
- ii. **Solve the diagonalized system. Remember that we only want a solution over the interval $t \in [i\Delta, (i + 1)\Delta)$. Use the value at $t = i\Delta$ as your initial condition.**
- iii. **Discretize the diagonalized system to obtain $\vec{y}_d[i]$. Show that**

$$\vec{y}_d[i + 1] = \underbrace{\begin{bmatrix} e^{\lambda_1\Delta} & 0 & \dots & 0 \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & e^{\lambda_n\Delta} \end{bmatrix}}_{e^{\Lambda\Delta}} \vec{y}_d[i] + \begin{bmatrix} \frac{e^{\lambda_1\Delta} - 1}{\lambda_1} & 0 & \dots & 0 \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & \frac{e^{\lambda_n\Delta} - 1}{\lambda_n} \end{bmatrix} \vec{u}_d[i] \quad (17)$$

Then, show that the matrix $\begin{bmatrix} \frac{e^{\lambda_1 \Delta} - 1}{\lambda_1} & 0 & \dots & 0 \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & \frac{e^{\lambda_n \Delta} - 1}{\lambda_n} \end{bmatrix}$ can be compactly written as $\Lambda^{-1}(e^{\Lambda \Delta} - I)$.

- iv. Undo the change of variables on the discretized diagonal system to get the discretized solution of the original system.

Solution:

- i. First, following the hint, we notice that with a full set of distinct eigenvalues and corresponding eigenvectors, we can change coordinates so that $\vec{x}(t) = V\vec{y}(t)$ and $\vec{y}(t) = V^{-1}\vec{x}(t)$. Using this transformation we diagonalize the system of differential equations, i.e

$$\frac{d\vec{x}(t)}{dt} = A\vec{x}(t) + \vec{b}u(t) \quad (18)$$

$$\implies \frac{dV\vec{y}(t)}{dt} = AV\vec{y}(t) + \vec{b}u(t) \quad (19)$$

$$\therefore \frac{d\vec{y}(t)}{dt} = V^{-1}AV\vec{y}(t) + V^{-1}\vec{b}u(t) \quad (20)$$

Note that using the basis of eigenvectors V , we've diagonalized A to get $\Lambda = \begin{bmatrix} \lambda_1 & 0 & \dots & 0 \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & \lambda_n \end{bmatrix}$

$$\therefore \frac{d\vec{y}(t)}{dt} = \Lambda\vec{y}(t) + V^{-1}\vec{b}u(t) \quad (21)$$

- ii. Now, we can use the fact that we care about the solution for $\vec{y}(t)$ over the interval $t \in (i\Delta, (i+1)\Delta]$, so $u(t)$ is a constant. Thus, we can write eq. (21) as follows:

$$\frac{d\vec{y}(t)}{dt} = \Lambda\vec{y}(t) + \underbrace{V^{-1}\vec{b}u_d[i]}_{\vec{u}_d[i]} \quad (22)$$

Notice that this system is diagonal (and hence we can write it as a system of n differential equations). We can look at the k th differential equation. We will use the subscripting notation $(\vec{y}(t))_k$ and $(\vec{u}_d[i])_k$ to denote the k th element of $\vec{y}(t)$ and $\vec{u}_d[i]$ respectively:

$$\frac{d(\vec{y}(t))_k}{dt} = \lambda_k(\vec{y}(t))_k + (\vec{u}_d[i])_k \quad (23)$$

We can pattern match to the solution in eq. (7), setting $\lambda \rightarrow \lambda_k$, $u_d[i] \rightarrow (\vec{u}_d[i])_k$, $b \rightarrow 1$, and $x(t) \rightarrow (\vec{y}(t))_k$, to get

$$(\vec{y}(t))_k = e^{\lambda_k(t-i\Delta)}(\vec{y}(i\Delta))_k + (\vec{u}_d[i])_k \int_{i\Delta}^t e^{\lambda_k(t-\theta)} d\theta \quad (24)$$

for $t \in (i\Delta, (i+1)\Delta]$.

iii. Now, we want to find $(\vec{y}_d[i+1])_k = (\vec{y}((i+1)\Delta))_k$, so we can plug in for $t = (i+1)\Delta$ in eq. (24) and we will get

$$(\vec{y}_d[i+1])_k = (\vec{y}((i+1)\Delta))_k = e^{\lambda_k \Delta} (\vec{y}(i\Delta))_k + \left(\frac{e^{\lambda_k \Delta} - 1}{\lambda_k} \right) (\vec{u}_d[i])_k \quad (25)$$

Since we have a solution for the k th differential equation in the system, we can arrange all the differential equations in this system in matrix form as follows:

$$\underbrace{\vec{y}((i+1)\Delta)}_{\vec{y}_d[i+1]} = \underbrace{\begin{bmatrix} e^{\lambda_1 \Delta} & 0 & \dots & 0 \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & e^{\lambda_n \Delta} \end{bmatrix}}_{e^{\Lambda \Delta}} \underbrace{\vec{y}(i\Delta)}_{\vec{y}_d[i]} + \begin{bmatrix} \frac{e^{\lambda_1 \Delta} - 1}{\lambda_1} & 0 & \dots & 0 \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & \frac{e^{\lambda_n \Delta} - 1}{\lambda_n} \end{bmatrix} \vec{u}_d[i] \quad (26)$$

Using the notation in the hint, we can write the second matrix in eq. (26) as:¹

$$\begin{bmatrix} \frac{e^{\lambda_1 \Delta} - 1}{\lambda_1} & 0 & \dots & 0 \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & \frac{e^{\lambda_n \Delta} - 1}{\lambda_n} \end{bmatrix} = \begin{bmatrix} \frac{e^{\lambda_1 \Delta}}{\lambda_1} & 0 & \dots & 0 \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & \frac{e^{\lambda_n \Delta}}{\lambda_n} \end{bmatrix} + \begin{bmatrix} \frac{-1}{\lambda_1} & 0 & \dots & 0 \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & \frac{-1}{\lambda_n} \end{bmatrix} \quad (27)$$

$$= \begin{bmatrix} \frac{1}{\lambda_1} & 0 & \dots & 0 \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & \frac{1}{\lambda_n} \end{bmatrix} \begin{bmatrix} e^{\lambda_1 \Delta} & 0 & \dots & 0 \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & e^{\lambda_n \Delta} \end{bmatrix} - \begin{bmatrix} \frac{1}{\lambda_1} & 0 & \dots & 0 \\ \vdots & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ 0 & \dots & \dots & \frac{1}{\lambda_n} \end{bmatrix} \quad (28)$$

$$= \Lambda^{-1} e^{\Lambda \Delta} - \Lambda^{-1} I \quad (29)$$

$$= \Lambda^{-1} (e^{\Lambda \Delta} - I) \quad (30)$$

This gives us

$$\vec{y}_d[i+1] = \vec{y}((i+1)\Delta) = e^{\Lambda \Delta} \underbrace{\vec{y}(i\Delta)}_{\vec{y}_d[i]} + \Lambda^{-1} (e^{\Lambda \Delta} - I) \vec{u}_d[i] \quad (31)$$

iv. Recall that $\vec{x}(t) = V\vec{y}(t)$ so $\vec{x}_d[i] = \vec{x}(i\Delta) = V\vec{y}(i\Delta) = V\vec{y}_d[i]$, and likewise, $\vec{y}_d[i] = V^{-1}\vec{x}_d[i]$. Using this form in the simplification, we find that:

$$\vec{x}_d[i+1] = V\vec{y}_d[i+1] \quad (32)$$

$$= V(e^{\Lambda \Delta} \vec{y}_d[i] + \Lambda^{-1} (e^{\Lambda \Delta} - I) \vec{u}_d[i]) \quad (33)$$

$$= (V e^{\Lambda \Delta} V^{-1}) \vec{x}_d[i] + (V \Lambda^{-1} (e^{\Lambda \Delta} - I)) \vec{u}_d[i] \quad (34)$$

Now, recall that our original goal was to write out A_d and \vec{b}_d , and we can do that now with our expression. Re-substituting $\vec{u}_d[i] = V^{-1}\vec{b}_u[i]$ we have:

$$\vec{x}_d[i+1] = (V e^{\Lambda \Delta} V^{-1}) \vec{x}_d[i] + (V \Lambda^{-1} (e^{\Lambda \Delta} - I)) V^{-1} \vec{b}_u[i] \quad (35)$$

¹In a matrix product, if both matrices are diagonal, the product is commutative.

$$= \underbrace{\left(V e^{\Lambda \Delta} V^{-1} \right)}_{A_d} \bar{x}_d[i] + \underbrace{\left(V \Lambda^{-1} \left(e^{\Lambda \Delta} - I \right) V^{-1} \vec{b} \right)}_{\vec{b}_d} u_d[i] \quad (36)$$

(c) Consider the discrete-time system

$$\bar{x}_d[i+1] = A_d \bar{x}_d[i] + \vec{b}_d u_d[i] \quad (37)$$

Suppose that $\bar{x}_d[0] = \bar{x}_0$. **Unroll the implicit recursion and show that the solution follows the form in eq. (38).**

$$\bar{x}_d[i] = A_d^i \bar{x}_d[0] + \left(\sum_{j=0}^{i-1} u_d[j] A_d^{i-1-j} \right) \vec{b}_d \quad (38)$$

You may want to verify that this guess works by checking the form of $\bar{x}_d[i+1]$. You don't need to worry about what A_d and \vec{b}_d actually are in terms of the original parameters.

(Hint: If we have a scalar difference equation, how would you solve the recurrence? Try writing $\bar{x}_d[i]$ in terms of $\bar{x}_d[0]$ for $i = 1, 2, 3$ and look for a pattern.)

Solution: Here, we derive the unrolled recursion and make a guess at the form of the solution in summation notation. Let's look at the pattern starting with $\bar{x}_d[1]$, given that $\bar{x}_d[i+1] = A_d \bar{x}_d[i] + \vec{b}_d u_d[i]$,

$$\bar{x}_d[1] = A_d \bar{x}_d[0] + \vec{b}_d u_d[0] \quad (39)$$

$$\bar{x}_d[2] = A_d \bar{x}_d[1] + \vec{b}_d u_d[1] \quad (40)$$

$$= A_d (A_d \bar{x}_d[0] + \vec{b}_d u_d[0]) + \vec{b}_d u_d[1] \quad (41)$$

$$= A_d^2 \bar{x}_d[0] + A_d \vec{b}_d u_d[0] + \vec{b}_d u_d[1] \quad (42)$$

$$\bar{x}_d[3] = A_d \bar{x}_d[2] + \vec{b}_d u_d[2] \quad (43)$$

$$= A_d \left(A_d^2 \bar{x}_d[0] + A_d \vec{b}_d u_d[0] + \vec{b}_d u_d[1] \right) + \vec{b}_d u_d[2] \quad (44)$$

$$= A_d^3 \bar{x}_d[0] + A_d^2 \vec{b}_d u_d[0] + A_d \vec{b}_d u_d[1] + \vec{b}_d u_d[2] \quad (45)$$

So, given this pattern, if we guess:

$$\bar{x}_d[i] = A_d^i \bar{x}_d[0] + \left(\sum_{j=0}^{i-1} u_d[j] A_d^{i-1-j} \right) \vec{b}_d \quad (46)$$

Then, let's see what we get for $\bar{x}_d[i+1]$, and make sure our guess is correct:

$$\bar{x}_d[i+1] = A_d \bar{x}_d[i] + \vec{b}_d u_d[i] \quad (47)$$

$$= A_d \left(A_d^i \bar{x}_d[0] + \left(\sum_{j=0}^{i-1} u_d[j] A_d^{i-1-j} \right) \vec{b}_d \right) + \vec{b}_d u_d[i] \quad (48)$$

$$= A_d^{i+1} \bar{x}_d[0] + \left(\left(\sum_{j=0}^{i-1} u_d[j] A_d^{i-j} \right) + u_d[i] \right) \vec{b}_d \quad (49)$$

$$= A_d^{i+1} \bar{x}_d[0] + \left(\sum_{j=0}^i u_d[j] A_d^{i-j} \right) \vec{b}_d \quad (50)$$

This satisfies (46), for $i+1$ and hence our guess was correct!

Contributors:

- Anish Muthali.

- Neelesh Ramachandran.
- Druv Pai.
- Anant Sahai.
- Nikhil Shinde.
- Sanjit Batra.
- Aditya Arun.
- Kuan-Yun Lee.
- Kumar Krishna Agrawal.