

# Discussion 9A

The following notes are useful for this discussion: [Note 13](#), [Note 15](#).

## 1. Towards Upper-Triangularization By An Orthonormal Basis

**Solution:** In lecture, we have been motivated by the goal of getting to a coordinate system in which the eigenvalues are on the diagonal, and there are only zeros below the diagonal. There can be "stuff" (nonzero entries) above the diagonal. When this is done to the  $A$  matrix representing a time-evolving system, we can view the system as a cascade of scalar systems — with each one potentially being an input to the ones that come "above" it.

Previously in this course, we have seen the value of changing our coordinates to be eigenbasis-aligned, because we can then view the system as a set of parallel scalar systems. Diagonalization causes these scalar equations to be fully uncoupled such that they can be solved separately. But even when we cannot diagonalize, we can *upper-triangularize* such that we can still solve the equations one at a time, from the "bottom up".

To better understand the steps involved, we will use the following concrete example:

$$M = S_{[3 \times 3]} = \begin{bmatrix} \frac{5}{12} & \frac{5}{12} & \frac{1}{6} \\ \frac{5}{12} & \frac{5}{12} & \frac{1}{6} \\ \frac{1}{6} & \frac{1}{6} & \frac{2}{3} \end{bmatrix} \quad (1)$$

and solve the general case by abstracting variables. Note that there is a datahub link to a jupyter notebook on the website, which will allow you to perform the numerical calculations quickly.<sup>1</sup>

- (a) Consider a non-zero vector  $\vec{u}_0 \in \mathbb{R}^n$ . Can you think of a way to extend it to a set of basis vectors for  $\mathbb{R}^n$ ? In other words, find  $\vec{u}_1, \dots, \vec{u}_{n-1}$ , such that  $\text{span}(\vec{u}_0, \vec{u}_1, \dots, \vec{u}_{n-1}) = \mathbb{R}^n$ . **To make things concrete, consider**  $\begin{bmatrix} 1 \\ -1 \\ 0 \end{bmatrix}$ . **Can you get an orthonormal basis where the first vector is a multiple of this vector?**

(HINT: What was the last discussion all about? Also, the given vector isn't normalized yet!)

**Solution:** Starting with the provided vector, we can include all the vectors from the standard basis of  $\mathbb{R}^3$ . Now, we know that the expanded matrix spans  $\mathbb{R}^n$  (since the 3 basis vectors alone do) and the initial vector can be treated as "extra" for now.) For a valid basis, we need a *minimal* set of spanning vectors; Gram-Schmidt will allow us to remove the redundancy.

For  $\begin{bmatrix} 1 & -1 & 0 \end{bmatrix}^\top$ , we can form:

$$\begin{bmatrix} | & | & | & | \\ \vec{v}_1 & \vec{v}_2 & \vec{v}_3 & \vec{v}_4 \\ | & | & | & | \end{bmatrix} = \begin{bmatrix} 1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad (2)$$

Then using this matrix (considering the constituent column vectors one at a time in order), we can run Gram-Schmidt to convert this matrix to an orthonormal basis (Gram-Schmidt will tell us which of the 4 vectors to throw out, leaving only 3). Namely, if we ever see a zero vector residual, we discard that vector and move on. A zero residual indicates that the current vector is in the span of the previous vectors in our final set.

The key concept is that we are guaranteed to span the whole space by the end. The standard basis alone does so, and Gram-Schmidt guarantees that the final span of our constructed vectors

<sup>1</sup>This particular matrix has an additional special property of symmetry, but we won't be invoking that here.

is the span of the input vectors. Starting with  $\vec{v}_1$  (using the same notation as [dis08B](#)):

$$\vec{q}_1 = \frac{\vec{v}_1}{\|\vec{v}_1\|} = \begin{bmatrix} \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} \\ 0 \end{bmatrix} = \begin{bmatrix} \frac{\sqrt{2}}{2} \\ -\frac{\sqrt{2}}{2} \\ 0 \end{bmatrix} \quad (3)$$

$$\Rightarrow \vec{z}_2 = \vec{v}_2 - (\vec{q}_1^\top \vec{v}_2) \vec{q}_1 \quad (4)$$

$$= \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} - \left( [1 \ 0 \ 0] \begin{bmatrix} \frac{\sqrt{2}}{2} \\ -\frac{\sqrt{2}}{2} \\ 0 \end{bmatrix} \right) \begin{bmatrix} \frac{\sqrt{2}}{2} \\ -\frac{\sqrt{2}}{2} \\ 0 \end{bmatrix} \quad (5)$$

$$= \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} - \frac{\sqrt{2}}{2} \begin{bmatrix} \frac{\sqrt{2}}{2} \\ -\frac{\sqrt{2}}{2} \\ 0 \end{bmatrix} \quad (6)$$

$$= \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} - \begin{bmatrix} \frac{1}{2} \\ -\frac{1}{2} \\ 0 \end{bmatrix} = \begin{bmatrix} \frac{1}{2} \\ \frac{1}{2} \\ 0 \end{bmatrix} \quad (7)$$

$$\Rightarrow \vec{q}_2 = \frac{\vec{z}_2}{\|\vec{z}_2\|} = \begin{bmatrix} \frac{\sqrt{2}}{2} \\ \frac{\sqrt{2}}{2} \\ 0 \end{bmatrix} \quad (8)$$

$$\vec{z}_3 = \vec{v}_3 - (\vec{q}_1^\top \vec{v}_3) \vec{q}_1 - (\vec{q}_2^\top \vec{v}_3) \vec{q}_2 = \vec{0} \quad (\text{unused in final basis}) \quad (9)$$

$$\vec{z}_4 = \vec{v}_4 - (\vec{q}_1^\top \vec{v}_4) \vec{q}_1 - (\vec{q}_2^\top \vec{v}_4) \vec{q}_2 \quad (10)$$

$$= \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} - 0 \begin{bmatrix} \frac{\sqrt{2}}{2} \\ -\frac{\sqrt{2}}{2} \\ 0 \end{bmatrix} - 0 \begin{bmatrix} \frac{1}{2} \\ \frac{1}{2} \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \quad (11)$$

$$\Rightarrow \vec{q}_3 = \frac{\vec{z}_4}{\|\vec{z}_4\|} \quad (12)$$

Therefore, our orthonormal matrix (consisting of the new basis vectors) is:

$$\begin{bmatrix} \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} & 0 \\ -\frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad (13)$$

- (b) Now consider a real eigenvalue  $\lambda_1$ , and the corresponding (normalized) eigenvector  $\vec{v}_1 \in \mathbb{R}^n$  of  $M \in \mathbb{R}^{n \times n}$  ( $M\vec{v}_1 = \lambda_1\vec{v}_1$ ). We know we can extend  $\vec{v}_1$  to an orthonormal basis of  $\mathbb{R}^n$ . We will denote the basis by

$$U = \begin{bmatrix} | & | & \cdots & | \\ \vec{u}_1 & \vec{u}_2 & \cdots & \vec{u}_n \\ | & | & \cdots & | \end{bmatrix} \quad (14)$$

where  $\vec{u}_1 = \vec{v}_1$  (note that this eigenvector is already normalized).

Our goal is to look at what the matrix  $M$  looks like in the coordinate system defined by the

basis  $U$ . **Compute  $U^\top M U$  by writing  $U = [\vec{v}_1 \ R]$ , where  $R := \begin{bmatrix} | & | & \cdots & | \\ \vec{r}_1 & \vec{r}_2 & \cdots & \vec{r}_{n-1} \\ | & | & \cdots & | \end{bmatrix}$ .** (Note :

$\vec{r}_i = \vec{u}_{i+1}$ )

**Solution:** Symbolic analysis:

$$U^\top M U = \begin{bmatrix} \vec{v}_1^\top \\ R^\top \end{bmatrix} M [\vec{v}_1 \ R] \quad (15)$$

$$= \begin{bmatrix} \vec{v}_1^\top \\ R^\top \end{bmatrix} [M\vec{v}_1 \quad MR] \quad (16)$$

$$= \begin{bmatrix} \vec{v}_1^\top \\ R^\top \end{bmatrix} [\lambda_1 \vec{v}_1 \quad MR] \quad (17)$$

$$= \begin{bmatrix} \lambda_1 \vec{v}_1^\top \vec{v}_1 & \vec{v}_1^\top MR \\ \lambda_1 R^\top \vec{v}_1 & R^\top MR \end{bmatrix}. \quad (18)$$

$$= \begin{bmatrix} \lambda_1 & \vec{v}_1^\top MR \\ \lambda_1 R^\top \vec{v}_1 & R^\top MR \end{bmatrix}. \quad (19)$$

Concrete case:  $S_{[3 \times 3]}$  has zero as eigenvalue since it contains a repeated column vector. So, we let the corresponding eigenvector be  $[1 \quad -1 \quad 0]^\top$ , the starting vector from the previous subpart. Note that this is in the nullspace of  $S_{[3 \times 3]}$ . Then, we have:

$$U = \begin{bmatrix} \frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} & 0 \\ -\frac{\sqrt{2}}{2} & \frac{\sqrt{2}}{2} & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad R = \begin{bmatrix} \frac{\sqrt{2}}{2} & 0 \\ \frac{\sqrt{2}}{2} & 0 \\ 0 & 1 \end{bmatrix} \quad (20)$$

Performing the matrix multiplication yields:

$$U^\top S_{[3 \times 3]} U = \begin{bmatrix} 0 & 0 & 0 \\ 0 & \frac{5}{6} & \frac{\sqrt{2}}{6} \\ 0 & \frac{\sqrt{2}}{6} & \frac{2}{3} \end{bmatrix} \quad (21)$$

From here, we can form a connection to the result of a couple subparts later, seeing that:

$$Q = R^\top S_{[3 \times 3]} R = \begin{bmatrix} \frac{5}{6} & \frac{\sqrt{2}}{6} \\ \frac{\sqrt{2}}{6} & \frac{2}{3} \end{bmatrix} \quad (22)$$

(c) **Verify that  $U^{-1} = U^\top$ , where  $U$  is the matrix we get from Gram-Schmidt process.**

**Solution:** One way to reason through this proof is with definitions and properties.  $U$  is an orthonormal basis by construction.  $U^\top U$  performs an inner product between each of the basis vectors. Since these basis vectors are orthogonal, all the non-diagonal elements have to be 0. Since the basis vectors are normalized, the inner product with itself is 1. As a result,  $U^\top U = I$ , and  $U^{-1} = U^\top$ .

We outline the same approach below in the  $3 \times 3$  case.

Suppose we have an orthonormal matrix  $P$ :

$$P = \begin{bmatrix} | & | & | \\ \vec{p}_1 & \vec{p}_2 & \vec{p}_3 \\ | & | & | \end{bmatrix} \quad (23)$$

We can compute  $P^\top P$ . We use the fact that for a set of mutually orthonormal vectors, the inner product of a vector with any *other* vector in the set is 0, but the inner product of a vector with itself is 1:

$$P^\top P = \begin{bmatrix} - & \vec{p}_1^\top & - \\ - & \vec{p}_2^\top & - \\ - & \vec{p}_3^\top & - \end{bmatrix} \begin{bmatrix} | & | & | \\ \vec{p}_1 & \vec{p}_2 & \vec{p}_3 \\ | & | & | \end{bmatrix} \quad (24)$$

$$= \begin{bmatrix} \vec{p}_1^\top \vec{p}_1 & \vec{p}_1^\top \vec{p}_2 & \vec{p}_1^\top \vec{p}_3 \\ \vec{p}_2^\top \vec{p}_1 & \vec{p}_2^\top \vec{p}_2 & \vec{p}_2^\top \vec{p}_3 \\ \vec{p}_3^\top \vec{p}_1 & \vec{p}_3^\top \vec{p}_2 & \vec{p}_3^\top \vec{p}_3 \end{bmatrix} \quad (25)$$

$$= \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad (26)$$

$$= I \quad (27)$$

This shows that  $P^\top = P^{-1}$ .

(d) **Look at the first column and the first row of  $U^\top MU$  and show that:**

$$M = U \begin{bmatrix} \lambda_1 & \vec{a}^\top \\ \vec{0} & Q \end{bmatrix} U^\top \quad (28)$$

where  $Q = R^\top MR$ . Here,  $\vec{a}$  is a vector related to  $M$ ,  $R$ , and  $\vec{v}_1$  (we will show the relation!).

**Solution:** We found above that:

$$U^\top MU = \begin{bmatrix} \lambda_1 & \vec{v}_1^\top MR \\ \lambda_1 R^\top \vec{v}_1 & R^\top MR \end{bmatrix} \quad (29)$$

Now, we need to show why:

$$\begin{bmatrix} \lambda_1 & \vec{v}_1^\top MR \\ \lambda_1 R^\top \vec{v}_1 & R^\top MR \end{bmatrix} \stackrel{?}{=} \begin{bmatrix} \lambda_1 & \vec{a}^\top \\ \vec{0} & Q \end{bmatrix} \quad (30)$$

We start simplifying the left side. First, we note that  $\lambda_1 R^\top \vec{v}_1 = \vec{0}$  because  $R$  consists of all of the other  $\vec{u}_i$  vectors that compose our orthonormal basis; taking the inner product between any one of these and  $\vec{v}_1$  yields zero (same logic as outlined in the previous part for vectors that are mutually orthogonal).

$\vec{v}_1^\top MR$  currently takes the place of  $\vec{a}^\top$ , suggesting that  $\vec{a} = (\vec{v}_1^\top MR)^\top = R^\top M^\top \vec{v}_1$ . So, finally substituting that  $Q = R^\top MR$ , we have:

$$U^\top MU = \begin{bmatrix} \lambda_1 & \vec{a}^\top \\ \vec{0} & Q \end{bmatrix} \quad (31)$$

We want an expression for  $M$  and so we can use the fact that  $U^\top = U^{-1}$  to see:

$$U^\top MU = \begin{bmatrix} \lambda_1 & \vec{a}^\top \\ \vec{0} & Q \end{bmatrix} \implies M = U \begin{bmatrix} \lambda_1 & \vec{a}^\top \\ \vec{0} & Q \end{bmatrix} U^\top \quad (32)$$

In the numerical example with  $S_{[3 \times 3]}$ , we have:

$$Q = R^\top S_{[3 \times 3]} R = \begin{bmatrix} \frac{5}{6} & \frac{\sqrt{2}}{6} \\ \frac{\sqrt{2}}{6} & \frac{2}{3} \end{bmatrix} \quad (33)$$

(e) Now, we can recurse on  $Q$  to get:

$$Q = [\vec{v}_2 \quad Y] \begin{bmatrix} \lambda_2 & \vec{b}^\top \\ \vec{0} & P \end{bmatrix} [\vec{v}_2 \quad Y]^\top \quad (34)$$

where we have taken  $\vec{v}_2 \in \mathbb{R}^{n-1}$ , a normalized eigenvector of  $Q$ , associated with eigenvalue  $\lambda_2$ . Again  $\vec{v}_2$  is extended into an orthonormal basis to form  $[\vec{v}_2 \quad Y]$ .

**Plug this form of  $Q$  into  $M$  above, to show that:**

$$M = [\vec{v}_1 \quad R\vec{v}_2 \quad RY] \begin{bmatrix} \lambda_1 & \check{a}_1 & \check{a}_{\text{rest}}^\top \\ 0 & \lambda_2 & \check{b}^\top \\ \vec{0} & \vec{0} & P \end{bmatrix} [\vec{v}_1 \quad R\vec{v}_2 \quad RY]^\top \quad (35)$$

where we define  $\check{\vec{a}}$  to be the "adjusted"  $\vec{a}$  to account for the substitution of  $Q$ ;  $\check{\vec{a}}^\top = \vec{a}^\top [\vec{v}_2 \ Y]$ .

**Solution:** From above, we know that

$$M = U \begin{bmatrix} \lambda_1 & \vec{a}^\top \\ \vec{0} & Q \end{bmatrix} U^\top \quad (36)$$

with  $U = [\vec{v}_1 \ R]$

In the given definition of  $Q$ , let's denote  $[\vec{v}_2 \ Y]$  as  $U_2$ , since this is the orthonormal basis that upper triangularizes  $Q$  (note the middle matrix of  $Q$ , which we can call  $T_2$ , is block upper-triangular). We can then write that:

$$Q = U_2 \underbrace{\begin{bmatrix} \lambda_2 & \vec{b}^\top \\ \vec{0} & P \end{bmatrix}}_{T_2} U_2^\top \quad (37)$$

We had an expression for  $Q$  previously;  $R^\top MR$ . We can equate the two representations and simplify:

$$U_2 T_2 U_2^\top = R^\top MR \quad (38)$$

$$T_2 = U_2^\top R^\top MR U_2 \quad (39)$$

$$= (R U_2)^\top MR U_2 \quad (40)$$

We know that  $T_2$  is an upper triangular matrix, so what the final equation above indicates is that the new orthonormal basis that upper triangularizes  $M$  *better than the original  $R$  basis*, is the  $R U_2$  basis. That is, instead of using  $[\vec{v}_1 \ R]$ , we want  $[\vec{v}_1 \ R U_2] = [\vec{v}_1 \ R \vec{v}_2 \ R Y]$ .

Motivated by this, we start with

$$M = U \begin{bmatrix} \lambda_1 & \vec{a}^\top \\ \vec{0} & U_2 T_2 U_2^\top \end{bmatrix} U^\top \quad (41)$$

$$= U \begin{bmatrix} 1 & \vec{0} \\ \vec{0} & U_2 \end{bmatrix} \begin{bmatrix} 1 & \vec{0} \\ \vec{0} & U_2^\top \end{bmatrix} \begin{bmatrix} \lambda_1 & \vec{a}^\top \\ \vec{0} & U_2 T_2 U_2^\top \end{bmatrix} \begin{bmatrix} 1 & \vec{0} \\ \vec{0} & U_2 \end{bmatrix} \begin{bmatrix} 1 & \vec{0} \\ \vec{0} & U_2^\top \end{bmatrix} U^\top \quad (42)$$

Here we have used the fact that  $U_2$  is an orthonormal matrix and that  $\begin{bmatrix} 1 & \vec{0} \\ \vec{0} & U_2 \end{bmatrix} \begin{bmatrix} 1 & \vec{0} \\ \vec{0} & U_2^\top \end{bmatrix} = I$

$$\therefore M = U \begin{bmatrix} 1 & \vec{0} \\ \vec{0} & U_2 \end{bmatrix} \begin{bmatrix} \lambda_1 & \vec{a}^\top U_2 \\ \vec{0} & U_2^\top U_2 T_2 U_2^\top U_2 \end{bmatrix} \begin{bmatrix} 1 & \vec{0} \\ \vec{0} & U_2^\top \end{bmatrix} U^\top \quad (43)$$

$$= [\vec{v}_1 \ R] \begin{bmatrix} 1 & \vec{0} \\ \vec{0} & U_2 \end{bmatrix} \begin{bmatrix} \lambda_1 & \vec{a}^\top U_2 \\ \vec{0} & T_2 \end{bmatrix} \begin{bmatrix} 1 & \vec{0} \\ \vec{0} & U_2^\top \end{bmatrix} [\vec{v}_1 \ R]^\top \quad (44)$$

$$= [\vec{v}_1 \ R U_2] \begin{bmatrix} \lambda_1 & \vec{a}^\top U_2 \\ \vec{0} & T_2 \end{bmatrix} [\vec{v}_1 \ R U_2]^\top \quad (45)$$

Defining the notation as  $\vec{a}^\top U_2 = \check{\vec{a}}^\top$ , we can finally write that:

$$M = [\vec{v}_1 \ R \vec{v}_2 \ R Y] \begin{bmatrix} \lambda_1 & \check{a}_1 & \check{\vec{a}}_{\text{rest}}^\top \\ 0 & \lambda_2 & \vec{b}^\top \\ \vec{0} & \vec{0} & P \end{bmatrix} [\vec{v}_1 \ R \vec{v}_2 \ R Y]^\top \quad (46)$$

We can be precise and write that  $\check{\vec{a}}_{\text{rest}}^\top = \check{\vec{a}}_{2:n-1}^\top$ .

The numerical results are:

$$Q = \begin{bmatrix} \frac{\sqrt{3}}{3} & \frac{\sqrt{6}}{3} \\ -\frac{\sqrt{6}}{3} & \frac{\sqrt{3}}{3} \end{bmatrix} \begin{bmatrix} \frac{1}{2} & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{\sqrt{3}}{3} & \frac{\sqrt{6}}{3} \\ -\frac{\sqrt{6}}{3} & \frac{\sqrt{3}}{3} \end{bmatrix}^\top \quad (47)$$

$$M = \begin{bmatrix} \frac{\sqrt{2}}{2} & -\frac{\sqrt{6}}{6} & \frac{\sqrt{3}}{3} \\ -\frac{\sqrt{2}}{2} & -\frac{\sqrt{6}}{6} & \frac{\sqrt{3}}{3} \\ 0 & \frac{\sqrt{6}}{3} & \frac{\sqrt{3}}{3} \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0.5 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{\sqrt{2}}{2} & -\frac{\sqrt{6}}{6} & \frac{\sqrt{3}}{3} \\ -\frac{\sqrt{2}}{2} & -\frac{\sqrt{6}}{6} & \frac{\sqrt{3}}{3} \\ 0 & \frac{\sqrt{6}}{3} & \frac{\sqrt{3}}{3} \end{bmatrix}^\top \quad (48)$$

(f) **(PRACTICE)** Show that the matrix  $[\vec{v}_1 \ R\vec{v}_2 \ RY]$  is still orthonormal.

**Solution:** To show that the matrix  $A = [\vec{v}_1 \ R\vec{v}_2 \ RY]$  is orthonormal, we want to show that the columns are mutually orthogonal, and all columns are unit vectors.

**Orthogonality:** We originally constructed the columns of  $R$  to be orthogonal to  $\vec{v}_1$ , as they were produced by the Gram-Schmidt algorithm. Thus  $\vec{v}_1^\top R\vec{v}_2 = 0$  and  $\vec{v}_1^\top RY = \vec{0}^\top$  since  $\vec{v}_1^\top R = \vec{0}^\top$ . As for the orthogonality of  $R\vec{v}_2$  and  $RY$ , we can see that

$$(R\vec{v}_2)^\top RY = \vec{v}_2^\top R^\top RY = \vec{v}_2^\top Y = \vec{0}^\top \quad (49)$$

for the reason that  $\vec{v}_2$  and the columns of  $Y$  were constructed to be orthogonal.

**Normality:** To check for normality (i.e all vectors are unit length), we can consider the inner products of each element with itself:

$$\vec{v}_1^\top \vec{v}_1 = 1 \quad (50)$$

$$(R\vec{v}_2)^\top R\vec{v}_2 = \vec{v}_2^\top R^\top R\vec{v}_2 \quad (51)$$

$$= \vec{v}_2^\top \vec{v}_2 = 1 \quad (52)$$

$$(RY)^\top RY = Y^\top R^\top RY \quad (53)$$

$$= Y^\top Y = I. \quad (54)$$

Note that the final calculation also assures us that  $RY$  has orthonormal columns.

(g) **(PRACTICE)** We have shown how to upper triangularize a  $3 \times 3$  and a  $2 \times 2$  matrix. **How can we generalize this process to any  $n \times n$  matrix  $M$ ?**

**Solution:**

In class, we've seen a recursive algorithm for upper-triangularization

---

#### Algorithm 1 UpperTriangularize

---

**Require:** matrix  $M$

- 1: **if**  $\dim(M) == 1$  **then**
  - 2:     **return**  $[1]$
  - 3: **else**
  - 4:      $\vec{v}_1 = \text{eigenvector}(M)$
  - 5:      $R = \text{GramSchmidtRest}(\vec{v}_1)$       $\triangleright$  Create the rest of an orthonormal matrix given  $\vec{v}_1$
  - 6:     Compute  $B = R^\top MR$       $\triangleright (n-1) \times (n-1)$  matrix
  - 7:      $U^1 = \text{UpperTriangularize}(B)$
  - 8:      $U = [\vec{v}_1 \ RU^1]$
  - 9:     **return**  $(U)$
  - 10: **end if**
- 

For any  $n \times n$  matrix  $M = M_n$ , we can decompose it into:

$$M_n = [\vec{v}_1 \ R_n] \begin{bmatrix} \lambda_1 & \vec{a}^\top \\ \vec{0} & M_{n-1} \end{bmatrix} [\vec{v}_1 \ R_n]^\top \quad (55)$$

$$= U_n A_n U_n^\top, \quad (56)$$

where  $M_{n-1}$  is an  $(n-1) \times (n-1)$  matrix.

We can recursively repeat this process on the submatrices  $M_i$  finding corresponding the  $U_i$ 's until we've reached the  $M_2$ , the  $2 \times 2$  case. Then we can combine these transformations from the bottom up, just like we did for the  $3 \times 3$  case, until we construct our final basis  $U_{n,\text{final}}$ :

$$U_{i,\text{final}} = [\vec{v}_{n-i+1} \quad R_i U_{i-1,\text{final}}] \quad (57)$$

Further, here in this part we see something more. Namely that we can actually do this in a single loop — the recursion can be transformed into a tail recursion. The key is that we can advance to get the the next vector in the basis directly – it is  $R\vec{v}_2$ .

---

### Algorithm 2 UpperTriangularizeLoop

---

**Require:** matrix  $M$

```

1: CurrentMatrix = M
2:  $U = []$  ▷ Need a place to accumulate the result
3:  $R = \text{Identity}(M)$  ▷ Same dimension as M to start, but this will accumulate the transformation
4: while  $\text{dim}(\text{CurrentMatrix}) > 0$  do
5:    $\vec{v} = \text{eigenvector}(\text{CurrentMatrix})$  ▷ Get one that is normalized
6:    $U = \text{columnstack}(U, R\vec{v})$  ▷ Add the new vector to the basis
7:   if  $\text{dim}(\text{CurrentMatrix}) == 1$  then
8:     CurrentMatrix =  $[]$ 
9:   else
10:     $Y = \text{GramSchmidtRest}(\vec{v})$  ▷ Create an orthonormal matrix given  $\vec{v}$ 
11:    CurrentMatrix =  $Y^\top \text{CurrentMatrix} Y$  ▷ One smaller than before
12:     $R = RY$  ▷ Update translation to original coordinates
13:   end if
14: end while
15: return ( $U$ )

```

---

Once we have our final basis  $U = U_{n,\text{final}}$ , we can transform into  $M$  into this basis to get our upper-triangular matrix  $T$ :

$$M = UTU^\top \quad (58)$$

$$T = U^\top MU. \quad (59)$$

- (h) **(PRACTICE) Show that the characteristic polynomial of square matrix  $M$  is the same as that of the square matrix  $UMU^{-1}$  for any invertible  $U$ .** You should use the key property  $\det(AB) = \det(A)\det(B)$  for square matrices.

**Solution:** The characteristic polynomial of the matrix  $M$  is given by  $\det(M - \lambda I)$ . Similarly the characteristic polynomial of  $UMU^{-1}$  is given by  $\det(UMU^{-1} - \lambda I)$ . Thus

$$\det(UMU^{-1} - \lambda I) = \det(UMU^{-1} - \lambda UU^{-1}) \quad (60)$$

$$= \det(U(M - \lambda I)U^{-1}) \quad (61)$$

$$= \det(U)\det(M - \lambda I)\det(U^{-1}). \quad (62)$$

Recognizing that  $\det(U) \cdot \det(U^{-1}) = 1$  we can simplify eq. (62) to:

$$\implies \det(UMU^{-1} - \lambda I) = \det(M - \lambda I). \quad (63)$$

Thus the characteristic polynomials of  $M$  and  $UMU^{-1}$  are the same for square matrices  $M$  and  $U$  where  $U$  is invertible.

#### Contributors:

- Neelesh Ramachandran.
- Yuxun Zhou.
- Edward Wang.
- Anant Sahai.
- Sanjit Batra.
- Pavan Bhargava.
- Kumar Krishna Agrawal.