

# Discussion 10B

## 1. Orthonormality and Least Squares

Recall that, if  $U \in \mathbb{R}^{m \times n}$  is a tall matrix (i.e.  $m \geq n$ ) with orthonormal columns, then

$$U^T U = I_{n \times n} \quad (1)$$

However, it is not necessarily true that  $U U^T = I_{m \times m}$ . In this discussion, we will deal with “orthonormal” matrices, where the term “orthonormal” refers to a matrix that is square with orthonormal columns and rows. Furthermore, for an orthonormal matrix  $U$ ,

$$U^T U = U U^T = I_{n \times n} \implies U^{-1} = U^T \quad (2)$$

This discussion will cover some useful properties that make orthonormal matrices favorable, and we will see a “nice” matrix factorization that leverages orthonormal matrices and helps us speed up least squares.

- (a) Suppose you have a real, square,  $n \times n$  orthonormal matrix  $U$ . You also have real vectors  $\vec{x}_1, \vec{x}_2, \vec{y}_1, \vec{y}_2$  such that

$$\vec{y}_1 = U \vec{x}_1 \quad (3)$$

$$\vec{y}_2 = U \vec{x}_2 \quad (4)$$

This is analogous to a change of basis. Show that, in this new basis, the inner products are preserved. Calculate  $\langle \vec{y}_1, \vec{y}_2 \rangle = \vec{y}_2^T \vec{y}_1 = \vec{y}_1^T \vec{y}_2$  in terms of  $\langle \vec{x}_1, \vec{x}_2 \rangle = \vec{x}_2^T \vec{x}_1 = \vec{x}_1^T \vec{x}_2$ .

- (b) Using the change of basis defined in part 1.a, show that, in the new basis, the norms are preserved. Express  $\|\vec{y}_1\|^2$  and  $\|\vec{y}_2\|^2$  in terms of  $\|\vec{x}_1\|^2$  and  $\|\vec{x}_2\|^2$ .

- (c) Suppose you observe data coming from the model  $y_i = \vec{a}^\top \vec{x}_i$ , and you want to find the linear scale-parameters (each  $a_i$ ). We are trying to learn the model  $\vec{a}$ . You have  $m$  data points  $(\vec{x}_i, y_i)$ , with each  $\vec{x}_i \in \mathbb{R}^n$ . Each  $\vec{x}_i$  is a different input vector that you take the inner product of with  $\vec{a}$ , giving a scalar  $y_i$ .

**Set up a matrix-vector equation of the form  $X\vec{a} = \vec{y}$  for some  $X$  and  $\vec{y}$ , and propose a way to estimate  $\vec{a}$ .**

- (d) Let's suppose that we can write our  $X$  matrix from part 1.c as

$$X = MV^\top \tag{5}$$

for some matrix  $M \in \mathbb{R}^{m \times n}$  and some orthonormal matrix  $V \in \mathbb{R}^{n \times n}$ . **Find an expression for  $\hat{\vec{a}}$  from the previous part, in terms of  $M$  and  $V^\top$ .**

Note: take this form as a given. We will go over how to find such a  $V$  and  $M$  later.

(e) Now suppose that we have the matrix

$$\begin{bmatrix} \vec{x}_1^\top \\ \vec{x}_2^\top \\ \vdots \\ \vec{x}_m^\top \end{bmatrix} := X = U\Sigma V^\top. \quad (6)$$

where  $U \in \mathbb{R}^{m \times m}$  is an orthonormal matrix, and  $V \in \mathbb{R}^{n \times n}$  is an orthonormal matrix. Here,

$$\Sigma = \begin{bmatrix} \sigma_1 & 0 & \dots & 0 \\ 0 & \sigma_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & \sigma_n \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{bmatrix}. \text{ Here we assume that we have more data points than the dimension of}$$

our space (that is,  $m > n$ ). Also, the transformation  $V$  in part e) is the same  $V$  in this factorized representation.

**Set up a least squares formulation for estimating  $\vec{a}$  and find the solution to the least squares.** Why might this factorization help us compute  $\hat{\vec{a}}$  faster?

Note: again, take this factorization as a given. We will go over how to find  $U$ ,  $\Sigma$ , and  $V$  later.

**Contributors:**

- Neelesh Ramachandran.
- Kuan-Yun Lee.
- Anant Sahai.
- Kumar Krishna Agrawal.